

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/535	A1	(11) International Publication Number: WO 99/58130 (43) International Publication Date: 18 November 1999 (18.11.99)
(21) International Application Number: PCT/US99/06523 (22) International Filing Date: 14 April 1999 (14.04.99) (30) Priority Data: 60/084,860 8 May 1998 (08.05.98) US (71) Applicant (for all designated States except US): PHARMACIA & UPJOHN COMPANY [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): TAYLOR, Duncan, Paul [US/US]; 8722 West F Avenue, Kalamazoo, MI 49009 (US). (74) Agent: WOOTTON, Thomas, A.; Intellectual Property Legal Services, Pharmacia & Upjohn Company, 301 Henrietta Street, Kalamazoo, MI 49001 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: NEW DRUG COMBINATIONS OF A N.A.R.I., PREFERABLY REBOXETINE, AND PINDOLOL (57) Abstract <p>This patent application describes a new combination treatment of selective, noradrenaline-reuptake inhibitors (NARI) and specifically, reboxetine, and pindolol to provide rapid relief to patients suffering from depression, general anxiety, attention deficit hyperactivity disorder (ADHD), anxiety disorders such as obsessive compulsive disorders (OCD), panic disorders (PD), social phobia (SD) and the like.</p> <p style="text-align: center;">BEST AVAILABLE COPY</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LJ	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

NEW DRUG COMBINATIONS OF A N.A.R.I., PREFERABLY REBOXETINE, AND PINDOLOL

Field of the Invention

5 This invention describes new treatments that should provide for a fast acting rapid onset of relief from several nervous system disorders, and it involves the administration of the drug reboxetine in combination with the drug pindolol.

Background

10 The introduction of tricyclic antidepressants in the early 1960s has provided a major advance in the treatment of neuropsychiatric disorders. Reactive and endogenous depressions, diagnoses formerly carrying grave prognostic implications, have become, with the introduction of the tricyclics, manageable disorders with a much smaller toll on the patient and the society as a whole.

15 The early tricyclic compounds were reuptake inhibitors of all the catecholamines released in the synaptic cleft, thus resulting in prolongation and enhancement of the dopamine (DA), noradrenaline (NA) and serotonin (5-hydroxytryptamine = 5-HT) action. Lack of selectivity also causes undesired side effects particularly on the acetylcholine (especially the muscarinic component), and histamine mediated neurotransmission.

20 Because of these unwanted pharmacodynamic activities, cognitive impairment, sedation, urinary and gastrointestinal tract disturbances, increased intraocular pressure were limiting factors in the clinical use of these compounds and often required discontinuation of treatment. Of utmost concern were also the cardiac toxic effects and the proconvulsant activity of this group of drugs.

25 More recently, selective reuptake inhibitors for serotonin (SSRI) have been introduced with definite advantages in regard to fewer side effects without loss of efficacy.

Summary of the Invention

30 Here we present the surprising finding that when the drug pindolol is given to a patient concurrently with a drug from a new category of antidepressants, a so called noradrenaline (NA) reuptake inhibitor (NARI), the combination of drugs act with surprising speed in relieving the symptoms of depression and it may be used for treating the symptoms of other central nervous system disorders including, but not only, general anxiety. Addictive Disorders, attention deficit hyperactivity disorder (ADHD), anxiety disorders such as

obsessive compulsive disorders (OCD), panic disorders (PD), social phobia (SP) and the like.

One particular NARI that is preferred is reboxetine. Reboxetine is the generic name of the pharmaceutical substance with the chemical name of 2-(1-((2-ethoxyphenoxy)benzyl)-morpholine, and its pharmaceutically acceptable salts. Reboxetine can be a free base, or it can include reboxetine methanesulfonate (also called reboxetine mesylate) or any other pharmaceutically acceptable salt that does not significantly affect the pharmaceutical activity of the substance.

The chemical name of pindolol is 1-(1H-Indol-4-yloxy)-3-[(1-methylethyl)amino]-2-propanol; 4-[2-hydroxy-3-(isopropylamino)-propoxy]indole; pinodolol. Pindolol is described in US patent no. 3,471,515, incorporated by reference and process steps are described in Swiss patents 469,002 and 472,404, assigned to the Sandoz Company, now the Novartis company, all documents incorporated into this document by reference. It has the trade name VISKEN®.

The present invention provides for the dosing of both reboxetine and pindolol, concurrently. The dosages for reboxetine and pindolol can be measured separately. The two drugs can be given as a single combined dose or given separately. They may be given at the same or at different times as long as both drugs are in the patient at one time over a 24 hour period. The two drugs will preferably be given to the patient, concomitantly, concurrently, at or about the same time, within about 5, 10, or 30 minutes, or they may be given within 1, 2, 3, 4, 5, 6, 8, 10, 12, 18 or about 24 hours, or fractions of minutes or of hours of each other. Concomitant or concurrent administration means the patient takes one drug within about 5 minutes of taking the other drug. Because the goal here is to provide rapid symptomatic relief to the patient, in most cases when treatment is started the two drugs would be administered to the patient close in time and typically concomitantly; thereafter, the timing of each drug's administration may not be as important.

A preferred dose range of reboxetine is 4 to 10 mg per patient per day and the more preferred dose is 6 to 8 mg or 8 to 10 mg per patient daily, depending upon the patient, delivered twice a day (b.i.d.). The reboxetine should be given to a patient concurrently with pindolol.

A preferred dose range of pindolol is 10-60 mg per patient per day and the more preferred dose is about 10 mg per patient daily, depending upon the patient, delivered twice

a day (b.i.d.). Preferably the pindolol should be given concurrently with reboxetine as described above.

5 **Additional Description of the Invention and Description of the Preferred Embodiment(s)**

Reboxetine is the generic name of the pharmaceutical substance with the chemical name of 2-(1-((2-ethoxyphenoxy)benzyl)-morpholine, and its pharmaceutically acceptable salts. Reboxetine can be a free base, or it can include reboxetine methanesulfonate (also
10 called reboxetine mesylate) or any other pharmaceutically acceptable salt that does not significantly affect the pharmaceutical activity of the substance. Reboxetine and a method of synthesis are described in U.S. 4,229,449, issued 21 Oct. 1980, Melloni et. al., incorporated by reference into this document, methods of preparation are described in US 5,068,433,
issued 26 Nov. 1991, Melloni et. al. and in US 5,391,735, issued 21 Feb. 1995, both
15 incorporated by reference. Reboxetine may also be known under the trade name of EDRONAX™.

The pharmaceutical compositions and methods of administration described in US 4,229,449 at col. 18, lines 33-66 are specifically incorporated by reference. Twice a day dosing is preferred with current formulations.

20 Reboxetine acts as an antidepressant. Antidepressants are frequently grouped into categories or "generations." The first generation of antidepressants were usually tricyclic antidepressants such as maprotiline that affected various neurotransmitter systems and are associated with many undesirable side effects. The second generation of antidepressants, such as mianserine, mirtazapine and trazodone are largely devoid of anticholinergic action
25 and their adrenolytic and antihistaminic effects are weaker. These are contrasted with the third generation of antidepressants (e.g. SSRI, ipsapirone, viloxazine, reboxetine, bupropione) that mediate only one of the three main neurotransmitter system for depression (5-HT, noradrenaline, dopamine) and they do not affect muscarine, histamine and adrenergic cerebral systems. Svestka, J. "Antidepressives of the 3rd, 4th and 5th generation," Cesk-
30 Psychiatr. 1994 Feb; 90(1):3-19. (Czech).

Reboxetine, however, does not act like most antidepressants. Unlike tricyclic antidepressants and even selective serotonin reuptake inhibitors (SSRIs), reboxetine is ineffective in the 8-OH-DPAT hypothermia test, indicating that reboxetine is not a selective

serotonin reuptake inhibitor rather it is selective for the noradrenergic system. Thus, reboxetine is not an SSRI, rather it is considered a novel, selective, noradrenaline-reuptake inhibitor (NARI). Leonard-BE, "Noradrenaline in basic models of depression." *European-Neuropsychopharmacol.* 1997 Apr; 7 Suppl 1: S11-6; discussion S71-3. Unlike most
5 drugs, reboxetine is a highly selective norepinephrine uptake inhibitor, with only marginal serotonin and no dopamine uptake inhibitory activity. The compound displays only weak or no anti-cholinergic activity in different animal models and is devoid of monoamine oxidase (MAO) inhibitory activity.

Reboxetine is highly potent and fast acting. Our investigations indicate reboxetine
10 has potent antireserpine activity and combines the inhibitory properties of classical tricyclic antidepressants on the reuptake of noradrenaline with an ability to desensitize β -adrenergic receptor function without showing any appreciable interaction with muscarinic cholinergic and α -adrenergic receptors. Moreover, reboxetine shows less vagolytic activity than other tricyclic antidepressants.

15 In spite of the inherently fast action of reboxetine there is still a "lag" or delay from the time of administration of the drug until the time the drug provides symptomatic relief to the patient. The treatments described here decrease that lag time. A period of days and especially weeks between the administration of a drug and its effect in relieving depression can be devastating to a patient. The patient may conclude the drug is not effective and stop
20 taking the drug, thus a quick onset of activity is critically important for treatments of this type. We have discovered that the combination of pinodolol and reboxetine provides highly effective relief of psychiatric disorders with a minimal delay in onset of activity.

Pindolol is the generic name for 1-(1H-Indol-4-yloxy)-3-[(1-methylethyl)amino]-2-propanol; 4-[2-hydroxy-3-(isopropylamino)-propoxy]indole; pinodolol. Pindolol is
25 described in US patent no. 3,471,515, incorporated by reference and process steps are described in Swiss patents 469,002 and 472,404, assigned to the Sandoz Company, now the Novartis company, all documents incorporated into this document by reference. It has the trade name VISKEN®.

The dosage used to treat all of the disorders described here may be found above and
30 below. Reboxetine is well tolerated and has a wide safety range; it can be administered in a dose range of active ingredient from about 1 to over 40 mg/kg. It is more commonly provided in dosages of from 1 to 20 mg per patient per day. Pindolol is also fairly safe although it is contraindicated for patients with bronchial asthmas, cardiac failure, heart block

and severe bradycardia. Other adverse reactions are possible. Pindolol dosages in the range of 5 to 60 mg daily can be effective. Both compounds may be administered by any suitable method including a convenient oral dosage form. A preferred method is oral dosing twice a day. The preferred dose range of reboxetine is 4 to 20 and more preferably 4 to 10 mg per patient per day and the preferred dose range of pindolol is 10 - 20 mg per patient per day. When starting medication the more preferred dose of reboxetine is 6 to 8 mg or 8 to 10 mg and pindolol is 10 mg per patient daily, depending upon the patient, delivered twice a day (b.i.d.). It can also be given at dosages of 2, 4, 6, 8, 10 or 12 mg/patient per day or fractions thereof. For example, suitable administrations could be 4 mg of reboxetine and 5 mg of pindolol in the morning and 2 or 4 mg of Reboxetine and 5 mg of pindolol in the evening. A skilled practitioner would be expected to determine the precise level of dosing. The idea dosing would be routinely determined by an evaluation of the patient and the needs of the patient.

This patent application describes the treatment of numerous conditions, disorders, diseases, and disease symptoms with the combination of drugs described herein, in addition to the conditions, disorders, diseases, and disease symptoms described above, the following may also be treated with these drugs: Addictive Disorders, Psychoactive Substance Use Disorders, Nicotine Addition or Tobacco Addiction (with a result of Smoking Cessation or a decrease in smoking) and Attention Deficit Hyperactivity Disorder (ADHD). This patent application also describes the treatment of Obsessive Compulsive Disorders (OCD), and Panic Disorder (PD), comprising administering a therapeutically effective, nontoxic dose of the drugs described herein and derivatives and or pharmaceutically acceptable salts thereof to a patient

Addictive Disorders and Psychoactive Substance Use Disorders, such as Intoxication disorders, Inhalation disorders, Alcohol addiction, Tobacco Addiction and or Nicotine Addiction. Tobacco and Nicotine addiction would be treated with the goal of achieving either Smoking Cessation or Smoking Reductions.

Addictive Disorders, Alcohol and Other Psychoactive Substance Use Disorders, disorders related to Intoxication and Inhalants and especially Tobacco Addiction or Nicotine Addiction, may be treated with the drugs described herein. Tobacco Addiction or Nicotine Addiction would be treated with the drugs described herein in order to achieve smoking/chewing cessation or smoking/chewing reduction. General descriptions of

Addictive Disorders, including disorders related to Intoxication and Inhalants and Tobacco Addiction or Nicotine Addiction may be found in many standard sources, such as, The American Psychiatric Press Textbook of Psychiatry, Second Edition, Edited by Robert E. Hales, Stuart C. Yudofsky, and John A. Talbott, copyright 1994, incorporated by reference, especially pp. 401 et. seq., section on "Nicotine" incorporated by reference. Another of many texts is the Manual of Psychiatric Therapeutics, Second Edition, edited by Richard I. Shader, incorporated by reference, especially pp. 85 from Chapter 11 (Hypnosis).

The treatment of Alcohol and Other Psychoactive Substance Use Disorders, such as disorders related to Intoxication and Inhalants and Tobacco Addiction or Nicotine Addiction but especially Tobacco Addiction involves the administration of the drugs described herein in a manner and form that provide a reduction in the symptoms of the disease. Tobacco Addiction or Nicotine Addiction in particular would be treated to achieve a reduction or cessation of smoking or chewing of nicotine containing materials by a patient. Cessation or a reduction in smoking or chewing of addictive or psychoactive substances involves the administration of the drugs described herein in a manner and form that provide a reduction in the symptoms of the disease, or with Tobacco or Nicotine with a reduction in the amount smoked or chewed.. See the general description above for administration of Reboxetine.

Attention Deficit Hyperactivity Disorder (ADHD)

ADHD is a condition or disease state that may be treated with the drugs described herein. General descriptions of ADHD, may be found in many standard sources, such as, The American Psychiatric Press Textbook of Psychiatry, Second Edition, Edited by Robert E. Hales, Stuart C. Yudofsky, and John A. Talbott, copyright 1994, incorporated by reference, especially pp. 741 et. al., section on "ADHD," incorporated by reference. Another of many texts is the Manual of Psychiatric Therapeutics, Second Edition, edited by Richard I. Shader, incorporated by reference, especially Chapter 18, Attention-Deficit hyperactivity Disorder, and pp. 172 et. seq., incorporated by reference.

The treatment of Attention Deficit Hyperactivity Disorder in children and adults involves the administration of the drugs described herein in a manner and form that provide a reduction in the symptoms of the disease. A child or young adult may require a smaller dosage depending upon the size, age, condition of the patient. See general description above for administration of the drugs described herein.

Obsessive Compulsive Disorders (OCD)

Obsessive Compulsive Disorder is a condition or state of anxiety that may be treated with reboxetine. General descriptions of OCD, may be found in many standard sources, such as, The American Psychiatric Press Textbook of Psychiatry, Second Edition, Edited by Robert E. Hales, Stuart C. Yudofsky, and John A. Talbott, copyright 1994, incorporated by reference, especially the chapter on "Anxiety Disorders," incorporated by reference. Another of many texts is the Manual of Psychiatric Therapeutics, Second Edition, edited by Richard I. Shader, incorporated by reference, especially Chapter 5, Obsessions and Compulsions, more particularly, Section III of that chapter, "OCD" pp. 36 *et. seq.*, incorporated by reference.

The treatment of Obsessive Compulsive Disorders (OCD) involves the administration of reboxetine in a manner and form that provide a reduction in the symptoms of the disease. See general description above for administration of reboxetine.

The following study shows the therapeutic effectiveness of using reboxetine in doses varying from 6 to 8 mg to treat OCD. This study is provided to illustrate the usefulness of using reboxetine as a treatment for OCD and the invention described herein should not be considered limited by this example.

In a trial involving 10 patients with a DSM-III-R diagnosis of Obsessive Compulsive Disorder who were all treated with reboxetine for a period of 3 to 4 weeks with the dose for the first week at 6 mg (4 mg in a.m. and 2 mg in p.m.) with the dose increasing in the second week to 8 mg (4 mg b.i.d.). At CGI last assessment, one patient was judged very much improved, 4 were judged much improved, 2 minimally improved, while 3 were unchanged. Of the patients who did respond they had a decrease of the obsessive-compulsive symptomatology, as measured by the CPRS-OC rating scale, of more than 30 and as much as 73%.

Panic Disorder (PD)

Panic Disorder is a condition or state of anxiety that may be treated with reboxetine. General descriptions of PD, may be found in many standard sources, such as, The American Psychiatric Press Textbook of Psychiatry, Second Edition, Edited by Robert E. Hales, Stuart C. Yudofsky, and John A. Talbott, copyright 1994, incorporated by reference, especially the chapter on "Anxiety Disorders," incorporated by reference, another of many texts is the Manual of Psychiatric Therapeutics, Second Edition, edited by Richard I. Shader, incorporated by reference, especially Chapter 25, "Approaches to the Treatment of Anxiety States," incorporated by reference.

The treatment of Panic Disorder involves the administration of the drugs described herein in a manner and form that provide a reduction in the symptoms of the disease. See general description above.

5

Claims

1. A single dosage form of reboxetine and pindolol.
 - 5 2. The single dose of claim 1 where the form is a tablet, hard or soft capsule or caplet.
 3. The single dose of claim 1 where the amount of reboxetine is 5 mg and the amount of pindolol is 7.5mg.
 - 10 4. A method of treating a patient experiencing symptoms selected from; depression, general anxiety disorders (GADs), Addictive Disorders, attention deficit hyperactivity disorder (ADHD), and anxiety disorders such as obsessive compulsive disorder (OCD), panic disorder (PD), social phobia (SP), comprising the administration of a therapeutically effective, nontoxic dose of pindolol, its derivatives and or pharmaceutically acceptable salts
15 thereof to a patient and administering a therapeutically effective, nontoxic dose of a selective, a noradrenaline reuptake inhibitor (NARI), its derivatives and or pharmaceutically acceptable salts thereof to a patient.
 5. A method of treating a patient as in claim 4 where the noradrenaline reuptake
20 inhibitor (NARI) is reboxetine.
 6. A method of treating a patient as in claim 5 where the administration of the dose of pindolol is administered within 24 hours of the administration of the dose of reboxetine.
 - 25 7. The method of claim 6 where the pindolol is administered within 12 hours of the reboxetine.
 8. The method of claim 7 where the pindolol is administered within 6 hours of the reboxetine.
30
 9. The method of claim 8 where the pindolol is administered within 3 hours of the reboxetine.
-

10. The method of claim 9 where the pindolol is administered within 1 hour of the reboxetine.
- 5 11. The method of claim 10 where the pindolol and reboxetine are administered concomitantly.
12. The method of claim 5 where the patient is experiencing symptoms of depression.
- 10 13. The method of claim 5 where the patient is experiencing symptoms of general anxiety disorders (GADs).
14. The method of claim 5 where the patient is experiencing symptoms of Addictive Disorders.
- 15 15. The method of claim 5 where the patient is experiencing symptoms of attention deficit hyperactivity disorder (ADHD).
16. The method of claim 5 where the patient is experiencing symptoms of obsessive
20 compulsive disorder (OCD).
17. The method of claim 5 where the patient is experiencing symptoms of panic disorder (PD).
- 25 18. The method of claim 5 where the patient is experiencing symptoms of social phobia (SP).
19. The use of reboxetine and pindolol, their derivatives and or pharmaceutically acceptable salts thereof in the manufacture of a medicament comprising an effective,
30 nontoxic dose of reboxetine and pindolol to treat general anxiety disorders (GADs), Addictive Disorders, attention deficit hyperactivity disorder (ADHD), and anxiety disorders such as obsessive compulsive disorder (OCD), panic disorder (PD), social phobia (SP),
and/or for the treatment of any of the symptoms of those diseases.

20. The method or use in claims 1-19 where the reboxetine dose range is from about 4 to 10 mg. per patient per day and the pindolol dose range is from about 10 to 20 mg. per patient per day, delivered twice a day.

5

21. The method or use in claims 1-19 where the reboxetine dose range is from about 6 to 8 mg. per patient per day and the pindolol dose range is from about 10 to 16 mg. per patient per day, delivered twice a day.

10

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/06523

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/535

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 687 472 A (LILLY CO ELI) 20 December 1995 (1995-12-20) page 2, line 28-32; claims 1,2,5,6,9,10,13-15 page 11, line 15-55 page 14, line 14-36 ---	1,4-12, 16,19
X	EP 0 759 299 A (LILLY CO ELI) 26 February 1997 (1997-02-26) page 12, line 57 - page 13, line 13; claims 1,2,4,5,7,9,13,14 --- -/-	1,2, 4-12,19

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

9 August 1999

Date of mailing of the international search report

19/08/1999

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kanbier, D

PCT/US 99/06523

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/06523

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 1-10
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 1-10
are directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☒ Claims Nos.: -
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
See FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/SA/ 210

Continuation of Box I.2

A compound cannot be sufficiently described by its mechanism of action or its pharmacological profile. A complete search can therefore not be made for "inhibitors" (claim 3).

The applicant's attention is drawn to the fact that claims relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. lional Application No

PCT/US 99/06523

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0687472 A	20-12-1995	AU 685510 B	22-01-1998
		AU 7742194 A	04-01-1996
		CA 2134038 A,C	17-12-1995
		CN 1113436 A	20-12-1995
		CZ 9402624 A	17-01-1996
		HU 71582 A	28-12-1995
		JP 8003035 A	09-01-1996
		NO 944046 A	18-12-1995
		NZ 264774 A	27-07-1997
		PL 305701 A	27-12-1995
		US 5532250 A	02-07-1996
		US 5552429 A	03-09-1996
		US 5538992 A	23-07-1996
		US 5532268 A	02-07-1996
		US 5532264 A	02-07-1996
		US 5532244 A	02-07-1996
		ZA 9408357 A	24-04-1996
EP 0759299 A	26-02-1997	AU 6776196 A	12-03-1997
		WO 9706792 A	27-02-1997

Form PCT/ISA/210 (patent family annex) (July 1992)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)